

## On Semicommutative Modules and Rings

NAZIM AGAYEV

*Abant İzzet Baysal University, Faculty of Science and Letters, Department of Mathematics, Gököy Campus, Bolu, Türkiye*

*e-mail* : agayev2005@yahoo.com

ABDULLAH HARMANCI

*Hacettepe University, Department of Mathematics, Ankara, Türkiye*

*e-mail* : harmanci@hacettepe.edu.tr

**ABSTRACT.** We say a module  $M_R$  a *semicommutative module* if for any  $m \in M$  and any  $a \in R$ ,  $ma = 0$  implies  $mRa = 0$ . This paper gives various properties of reduced, Armendariz, Baer, Quasi-Baer, *p.p.* and *p.q.*-Baer rings to extend to modules. In addition we also prove, for a *p.p.*-ring  $R$ ,  $R$  is semicommutative iff  $R$  is Armendariz. Let  $R$  be an abelian ring and  $M_R$  be a *p.p.*-module, then  $M_R$  is a semicommutative module iff  $M_R$  is an Armendariz module. For any ring  $R$ ,  $R$  is semicommutative iff  $A(R, \alpha)$  is semicommutative. Let  $R$  be a reduced ring, it is shown that for number  $n \geq 4$  and  $k = [n/2]$ ,  $T_n^k(R)$  is semicommutative ring but  $T_n^{k-1}(R)$  is not.

### 1. Introduction

Throughout this paper all rings  $R$  are associative with unity and all modules  $M$  are unital right  $R$ -modules. For a nonempty subset  $X$  of a ring  $R$ , we write  $r_R(X) = \{r \in R \mid Xr = 0\}$  and  $l_R(X) = \{r \in R \mid rX = 0\}$ , which are called the *right annihilator of  $X$*  in  $R$  and the *left annihilator of  $X$*  in  $R$ , respectively. The notation “ $\leq$ ” will denote a submodule. Recall that a ring  $R$  is *reduced* if  $R$  has no nonzero nilpotent elements. Observe that reduced rings are abelian (i.e., all idempotents are central). In [5] Kaplansky introduced *Baer* rings as rings in which the right (left) annihilator of every nonempty subset is generated by an idempotent. A ring  $R$  is called *quasi-Baer* if the right annihilator of each right ideal of  $R$  is generated (as a right ideal) by an idempotent. These definitions are left-right symmetric. A ring  $R$  is called a *right* (resp. *left*) *principally quasi-Baer* (or simply *right* (resp. *left*) *p.q.-Baer*) ring if the right (resp. left) annihilator of a principally right (resp. left) ideal of  $R$  is generated by an idempotent.  $R$  is called a *p.q.-Baer* ring if it is both right and left *p.q.-Baer*.

Another generalization of Baer rings is a *p.p.*-ring. A ring  $R$  is called a *right*

---

Received September 16, 2005.

2000 Mathematics Subject Classification: 16S36, 16W60, 16W99.

Key words and phrases: reduced rings (modules), Baer rings (modules) and quasi-Baer rings (modules) and semicommutative rings (modules).

(resp. *left*) *p.p.*-ring if the right (resp. left) annihilator of an element of  $R$  is generated by an idempotent.  $R$  is called a *p.p.*-ring if it is both a right and left *p.p.*-ring. A ring  $R$  is called *Armendariz* if whenever polynomials  $f(x) = \sum a_i x^i \in R[x], g(x) = \sum b_j x^j \in R[x]$  satisfy  $f(x)g(x) = 0$ , we have  $a_i b_j = 0$  for every  $i$  and  $j$ . A ring  $R$  is called *semicommutative* if for every  $a \in R$ ,  $r_R(a)$  is an ideal of  $R$ . (equivalently, for any  $a, b \in R$ ,  $ab = 0$  implies  $aRb = 0$ ). An idempotent  $e \in R$  is called *central* if  $xe = ex$  for all  $x \in R$ . An idempotent  $e^2 = e \in R$  is called a *left* (resp. *right*) *semicentral idempotent* if  $eR$  (resp.  $Re$ ) is a two sided ideal of  $R$ .

According to Lee-Zhou [6], a module  $M_R$  is called  $\alpha$ -*reduced* if, for any  $m \in M$  and any  $a \in R$ ,

- (1)  $ma = 0$  implies  $mR \cap Ma = 0$
- (2)  $ma = 0$  iff  $m\alpha(a) = 0$ ,

where  $\alpha : R \rightarrow R$  is a ring homomorphism with  $\alpha(1) = 1$ . The module  $M_R$  is called *reduced* if  $M_R$  is 1-reduced. In [8] Lee-Zhou introduced a Baer, quasi-Baer and *p.p.*- module as follows: (a)  $M_R$  is called *Baer* if, for any subset  $X$  of  $M$ ,  $r_R(X) = eR$  where  $e^2 = e \in R$ . (b)  $M_R$  is called *quasi-Baer* if, for any submodule  $N$  of  $M$ ,  $r_R(N) = eR$  where  $e^2 = e \in R$ . (c)  $M_R$  is called *p.p.* if, for any  $m \in M$ ,  $r_R(m) = eR$  where  $e^2 = e \in R$ . In [2] the module  $M_R$  is called *principally quasi-Baer* (*p.q.-Baer* for short) if, for any  $m \in M$ ,  $r_R(mR) = eR$  where  $e^2 = e \in R$ . In [3], the module  $M_R$  is *semicommutative* module if for any  $m \in M$  and any  $a \in R$ ,  $ma = 0$  implies  $mRa = 0$ , and the module  $M_R$  is called *Armendariz* if whenever polynomials  $m(x) = \sum m_i x^i \in M[x], f(x) = \sum a_j x^j \in R[x]$  satisfy  $m(x)f(x) = 0$ , we have  $m_i a_j = 0$  for every  $i$  and  $j$ .

Let  $M$  be a right  $R$ -module and  $S = \text{End}_R(M)$ . Then  $M$  is a left  $S$ -module, right  $R$ -module and  $S$ - $R$ -bimodule. In [11], Rizvi and Roman call  $M$  a *Baer module* if the right annihilator in  $M$  of any left ideal of  $S$  is generated by an idempotent of  $S$  (or equivalently, for all  $R$ -submodules  $N$  of  $M$ ,  $l_S(N) = Se$  with  $e^2 = e \in S$ ); and  $M$  is a *quasi-Baer* module if the right annihilator in  $M$  of any ideal of  $S$  is generated by an idempotent of  $S$  (or equivalently, for all fully invariant  $R$ -submodules  $N$  of  $M$ ,  $l_S(N) = Se$  with  $e^2 = e \in S$ ). Among others they have proved that any direct summand of a Baer (resp. quasi-Baer) module  $M$  is again a Baer (respect. quasi-Baer) module, and the endomorphism ring  $S = \text{End}_R(M)$  of a Baer (resp. quasi-Baer) module  $M$  is a Baer (resp. quasi-Baer) ring (see Theorem 4.1 in [11]). They gave several results for a direct sum of Baer (resp. quasi-Baer) modules to be a Baer (resp. quasi-Baer) module.

We shortly summarize the content of the paper. In [1, Proposition 2.7] it is shown that if  $M_R$  is a semicommutative module, then  $M_R$  is a Baer module if and only if it is a quasi-Baer module, and  $M_R$  is a *p.p.*-module if and only if it is a *p.q.-Baer* module. In Proposition 2.7 we prove that for an abelian ring  $R$  and a *p.p.*-module  $M_R$ ,  $M_R$  is a semicommutative module if and only if it is an Armendariz module. In Proposition 2.11 for a semicommutative ring  $R$  we show that  $R$  is a *p.p.*-ring if and only if  $R[x]$  is a *p.p.*-ring,  $R$  is a Baer ring if and only

if  $R[x]$  is a Baer ring,  $R$  is a p.q.-Baer ring if and only if  $R[x]$  is a p.q.-Baer ring, and  $R$  is a quasi-Baer ring if and only if  $R[x]$  is a quasi-Baer ring. In Proposition 2.13 we prove that for any ring  $R$ ,  $R$  is semicommutative if and only if  $A(R, \alpha)$  is semicommutative, and in Theorem 2.15 for a reduced ring  $R$  and any integer  $n \geq 4$  and  $k = \lceil n/2 \rceil$ , we show that  $T_n^k(R)$  is semicommutative ring but  $T_n^{k-1}(R)$  is not.

### Examples.

1. Every reduced module is semicommutative but the inverse is not true. For example,  $\mathbb{Z}_n$  is semicommutative for any  $n \in \mathbb{N}$ , but is reduced for only square-free  $n$ .

2. For any commutative ring  $R$ , any module  $M_R$  is semicommutative.

3. Let  $D$  be a division ring,  $R = \begin{bmatrix} D & D \\ 0 & D \end{bmatrix}$ , and  $A = \begin{bmatrix} 0 & D \\ 0 & D \end{bmatrix}$ . Then  $A_R$  is a semicommutative module.

From [4, Example 2] and [10, Proposition 4.6] we want to restate next results:

1. If a module  $M_R$  is semicommutative then  $M[x]_{R[x]}$  need not to be semicommutative.

2. If a semicommutative module  $M_R$  is Armendariz, then  $M[x]_{R[x]}$  is a semicommutative module.

## 2. Semicommutative modules

We start with some preliminary results on semicommutative modules and rings. Some of the results are known but we state and give their proofs for the sake of completeness.

**Lemma 2.1.** *Let  $M_R$  be a semicommutative module.*

(1) *If  $e^2 = e \in R$  with  $r_R(m) = eR$  for some  $m \in M$ , then  $e$  is left semicentral idempotent.*

(2) *For any  $e^2 = e \in R$ ,  $mea = mae$  for all  $m \in M$  and all  $a \in R$ .*

*Proof.* (1) Let  $e^2 = e \in R$  with  $r_R(m) = eR$  for some  $m \in M$ . Then  $me = 0$ . To prove  $e$  is left semicentral we show  $teR \leq eR$  for any  $t \in R$ . For any  $t \in R$ , then  $met = 0$ . By (1),  $mte = 0$ . Hence  $te \in r_R(m) = eR$ , and so  $teR \leq eR$ . This completes the proof.

(2) See also [1] for a proof. For  $e^2 = e \in R$ ,  $e(1-e) = (1-e)e = 0$ . Then for all  $m \in M$ ,  $me(1-e) = 0$  and  $m(1-e)e = 0$ . Since  $M_R$  is semicommutative, we have  $meR(1-e) = 0$  and  $m(1-e)Re = 0$ . Thus, for all  $a \in R$ ,  $mea(1-e) = 0$  and  $m(1-e)ae = 0$ . So,  $mea = meae$  and  $mae = meae$ . Hence,  $mea = mae$  for all  $a \in R$ .  $\square$

**Proposition 2.2.** *Let  $M$  be a semicommutative module. Then the following conditions are equivalent:*

(1)  *$M_R$  is a p.q.-Baer module.*

(2) *The right annihilator of every finitely generated submodule is generated (as a right ideal) by an idempotent.*

*Proof.* (1) $\Rightarrow$ (2) Assume that  $M_R$  is  $p.q.$ -Baer and  $N = \sum_{i=1}^k n_i R$  is a finitely generated submodule of  $M_R$ . Then  $r_R(N) = \bigcap_{i=1}^k e_i R$  where  $r_R(n_i R) = e_i R$  and  $e_i^2 = e_i$ . By Lemma 2.1 each  $e_i$  is a left semicentral idempotent, a routine argument yields an idempotent  $e$  such that  $\bigcap_{i=1}^k e_i R = eR$ . Therefore,  $r_R(N) = eR$ .  $\square$

**Corollary 2.3** [2, Prop. 1.7]. *Following conditions are equivalent for a ring  $R$ :*

- (1)  $R$  is a right  $p.q.$ -Baer ring.
- (2) The right annihilator of every finitely generated ideal of  $R$  is generated (as a right ideal) by an idempotent.

**Proposition 2.4.** *Let  $M_R$  be a semicommutative module. Consider the following properties:*

- (1)  $M_R$  is a Baer module.
- (2)  $M_R$  is a quasi-Baer module.
- (3)  $M_R$  is a  $p.p.$ -module.
- (4)  $M_R$  is a  $p.q.$ -Baer module.

*Then (1)  $\Leftrightarrow$  (2)  $\Rightarrow$  (3)  $\Leftrightarrow$  (4).*

*Proof.* See [1, Proposition 2.7].  $\square$

**Proposition 2.5.** *Let  $R$  be an abelian ring and  $M_R$  be a  $p.p.$  module. Then  $M_R$  is a  $p.q.$ -Baer module.*

*Proof.* See [1, Proposition 2.15].  $\square$

**Corollary 2.6.** *Abelian right  $p.p.$  rings are right  $p.q.$ -Baer.*

**Proposition 2.7.** *Let  $R$  be an abelian ring and  $M_R$  be a  $p.p.$ -module. Then following conditions are equivalent:*

- (1)  $M_R$  is a semicommutative module.
- (2)  $M_R$  is an Armendariz module.

*Proof.* (1) $\Rightarrow$ (2) Let  $m(x) = \sum m_i x^i \in M[x], f(x) = \sum a_j x^j \in R[x]$  satisfy  $m(x)f(x) = 0$ . We have the following system of equations:

$$\begin{aligned} m_0 a_0 &= 0 && \dots\dots\dots (1) \\ m_0 a_1 + m_1 a_0 &= 0 && \dots\dots\dots (2) \\ m_0 a_2 + m_1 a_1 + m_2 a_0 &= 0 && \dots\dots\dots (3) \\ &&& \vdots \\ m_t a_s &= 0 && \dots\dots\dots (n) \end{aligned}$$

for some  $s$  and  $t$ .

Let  $r(m_0) = e_0 R, r(m_1) = e_1 R, r(m_2) = e_2 R, \dots, r(m_t) = e_t R$  for some idempotents  $e_0, e_1, e_2, \dots, e_t$  in  $R$ . Then  $a_0 \in r(m_0) = e_0 R$ . Since  $R$  is abelian,  $a_0 = e_0 a_0 = a_0 e_0$ . Multiply (2) from right by  $e_0$  to obtain

$$m_0 a_1 e_0 + m_1 a_0 e_0 = 0$$

From semicommutativity of  $M_R$ ,  $m_0 e_0 = 0$  implies  $m_0 a_1 e_0 = 0$ . Then  $m_1 a_0 e_0 =$

$m_1a_0 = 0$ . So  $m_0a_1 = 0$  from (2).

Now multiply (3) from right by  $e_0$ :

$$m_0a_2e_0 + m_1a_1e_0 + m_2a_0e_0 = 0$$

From semicommutativity of  $M_R$ ,  $m_0e_0 = 0$  implies  $m_0a_2e_0 = 0$ . Then  $m_1a_1e_0 + m_2a_0e_0 = m_1a_1 + m_2a_0 = 0$

Hence  $m_0a_2 = 0$  by (3). Continuing this process we get:

$$m_0a_0 = m_0a_1 = m_0a_2 = \dots = m_0a_s = 0.$$

If we use these equalities, the equations (2), (3),  $\dots$ , will be :

$$\begin{aligned} m_1a_0 &= 0 && \dots\dots\dots (2') \\ m_1a_1 + m_2a_0 &= 0 && \dots\dots\dots (3') \\ &&& \vdots \\ m_t a_s &= 0 && \dots\dots\dots (n') \end{aligned}$$

Applying the same method to these equalities we get  $m_1a_0 = m_1a_1 = m_1a_2 = \dots = m_1a_s = 0$ .

Continuing this process we will have:

$$m_2a_0 = m_2a_1 = m_2a_2 = \dots = m_2a_s = 0 \dots m_t a_s = 0.$$

So  $m_i a_j = 0$  for any  $i, j$ .

(2) $\Rightarrow$ (1): Let  $ma = 0$  for  $m \in M, a \in R$ . For any  $r \in R$  take  $m(x) = mx + mr \in M[x]$  and  $f(x) = -ax + ra \in R[x]$ . Then  $m(x)f(x) = (mx + mr)(-ax + ra) = mr^2a$ . By hypothesis  $ma = 0$  implies  $a \in r(m) = eR$  for some idempotent  $e \in R$ . Then  $a = ae = ea$ . So,  $m(x)f(x) = mr^2a = mr^2ea = mer^2a = 0$  since  $R$  is abelian. As  $M_R$  is Armendariz, we get  $mra = 0$  for any  $r \in R$ .  $\square$

**Corollary 2.8.** *Let  $R$  be a p.p.-ring. Then the following are equivalent:*

- (1)  $R$  is a semicommutative ring.
- (2)  $R$  is an Armendariz ring.

*Proof.* In Lemma 2.1 take  $M = R$ , then every semicommutative ring is abelian for (1) $\Rightarrow$ (2) and from [6, Lemma 7] every Armendariz ring is abelian for (2) $\Rightarrow$ (1).  $\square$

**Proposition 2.9.** *Let  $M_R$  be a semicommutative module and  $R$  is a reduced module. Then  $M_R$  is an Armendariz module if and only if its torsion submodule  $T(M)$  is Armendariz.*

*Proof.* Assume that the torsion submodule  $T(M)$  of  $M$  is Armendariz as a right  $R$ -module. Let  $m(x) = \sum m_j x^j \in M[x], f(x) = \sum a_i x^i \in R[x]$  satisfy  $m(x)f(x) = 0$ . We have the following system of equations:

$$\begin{aligned} m_0a_0 &= 0 \\ m_0a_1 + m_1a_0 &= 0 \\ m_0a_2 + m_1a_1 + m_2a_0 &= 0 \\ &\vdots \\ m_t a_s &= 0 \end{aligned}$$

for some  $s$  and  $t$ .

We may assume  $a_0 \neq 0$ . Multiplying by  $a_0$ , the second of these equations yields  $m_1 a_0^2 = 0$ . Thus  $a_0^2$  annihilates both  $m_0$  and  $m_1$ . The third equation now implies  $m_2 a_0^3 = 0$ . Continuing we get  $m(x) \in T(M)[x]$ . Since  $T(M)$  is Armendariz as  $R$ -module, we conclude that  $m_j a_i = 0$  for all  $i, j$ . The other implication is trivial.  $\square$

Given a ring  $R$ , the formal power series ring over  $R$  is denoted by  $R[[x]]$ .

**Lemma 2.10.** *Let  $R$  be a semicommutative ring.*

- (1) *Every idempotent of  $R[x]$  is in  $R$ .*
- (2) *Every idempotent of  $R[[x]]$  is in  $R$ .*

*Proof.* From [6, Lemma 8].  $\square$

**Proposition 2.11.** *Let  $R$  be a semicommutative ring.*

- (1)  *$R$  is a p.p.-ring if and only if  $R[x]$  is a p.p.-ring.*
- (2)  *$R$  is a Baer ring if and only if  $R[x]$  is a Baer ring.*
- (3)  *$R$  is a p.q.-Baer ring if and only if  $R[x]$  is a p.q.-Baer ring.*
- (4)  *$R$  is a quasi-Baer ring if and only if  $R[x]$  is a quasi-Baer ring.*

*Proof.* (1) Assume that  $R$  is a p.p.-ring. From Corollary 2.16,  $R$  is an Armendariz ring. Then by [6, Theorem 9],  $R[x]$  is a p.p.-ring.

Conversely, assume that  $R[x]$  is a p.p.-ring. Let  $a \in R$ . By Lemma 2.10 there exists an idempotent  $e \in R$  such that  $r_{R[x]}(a) = eR[x]$ . Hence  $r_R(a) = r_{R[x]}(a) \cap R = eR$  and therefore  $R$  is a p.p.-ring.

(2) Assume that  $R$  is a Baer ring. Then  $R$  is a p.p.-ring. By Corollary 2.16  $R$  is an Armendariz ring. From [6, Theorem 10]  $R[x]$  is a Baer ring.

Conversely, assume that  $R[x]$  is a Baer ring. Let  $B$  be a nonempty subset of  $R$ . Then  $r_{R[x]}(B) = eR[x]$  for some idempotent  $e \in R$  by Lemma 2.10. Hence  $r_R(B) = eR$  and therefore  $R$  is a Baer ring.

(3) Assume that  $R$  is a p.q.-Baer ring. Let  $t(x) = a_0 + a_1x + \cdots + a_nx^n \in R[x]$ . By assumption  $r_R(a_i) = e_iR = r_R(a_iR)$ , for all  $i = 0, 1, 2, \dots, n$ . By Proposition 2.5  $\bigcap_{i=0}^n r_R(a_iR) = eR$ ,  $e = e_0e_1 \cdots e_n$ . Let  $f(x) \in r_{R[x]}(t(x)R[x])$ . Then  $t(x)R[x]f(x) = 0$  implies  $t(x)Rf(x) = 0$  and  $a_jRf(x) = 0$  for all  $j = 0, 1, 2, \dots, n$ . So  $a_jRb_i = 0$ , hence  $b_i \in \bigcap_{j=0}^n r_R(a_jR) = eR$  and  $b_i = eb_i$  for all  $i, j$ . Then  $ef(x) = f(x)$  implies  $f(x) \in eR[x]$ .

Conversely, assume that  $R[x]$  is a p.q.-Baer. Let  $a \in R$ . There exists idempotent  $e \in R$  such that  $r_{R[x]}(aR[x]) = eR[x]$ . Then  $r_{R[x]}(aR[x]) \cap R = (eR[x]) \cap R = eR$ . Since  $r_R(aR) = r_{R[x]}(aR[x]) \cap R$ , we get  $r_R(aR) = eR$ .

(4) Assume that  $R$  is a quasi-Baer. Let  $A$  be an ideal of  $R[x]$  and  $A^*$  be the set of all coefficients of elements of  $A$ . Then  $A^*$  is an ideal of  $R$ , so  $r_R(A^*) = eR$  for some idempotent  $e \in R$ . Since  $e \in r_{R[x]}(A)$ , we get  $eR[x] \subseteq r_{R[x]}(A)$ . Now, let  $f = b_0 + b_1x + \cdots + b_nx^n \in r_{R[x]}(A)$ . Then  $Af = 0$  implies  $Ab_i = 0$ , so  $A^*b_i = 0$  for all  $i = 0, 1, 2, \dots, n$ . Hence  $b_i \in r_R(A^*) = eR$  and  $b_i = eb_i$  for all  $i$ . Consequently,  $f \in eR[x]$ .

Conversely, assume that  $R[x]$  is a quasi-Baer ring and  $A$  is an ideal of  $R$ . Then  $A[x]$  is an ideal of  $R[x]$ . Hence  $r_{R[x]}(A[x]) = eR[x]$ . Intersecting both sides with  $R$  we get  $r_{R[x]}(A[x]) \cap R = eR[x] \cap R = eR$ . Since  $r_R(A) = r_{R[x]}(A[x]) \cap R$ , we have  $r_R(A) = eR$ .  $\square$

**Proposition 2.12.** *Let  $M_R$  be a p.p.-module. Then  $M_R$  is a semicommutative module if and only if  $mre = mer$ , for any  $m \in M, r \in R$ , and  $e^2 = e \in R$ .*

*Proof.* Assume that  $mre = mer$ , for any  $m \in M, r \in R$  and  $e^2 = e \in R$ . Let  $ma = 0$  for  $m \in M, a \in R$ . Then  $a \in r(m)$ . By hypothesis  $r(m) = eR$  for some  $e^2 = e \in R$ . Hence  $me = 0, a = ea$ , and so  $mra = mrea$ . By assumption  $mrea = mrae = mera = 0$  for any  $r \in R$ . So  $mra = 0$  for any  $r \in R$ . The rest is clear from Lemma 2.1.  $\square$

Now we consider D. A. Jordan's construction of the ring  $A(R, \alpha)$  (See [5] for more details). Let  $A(R, \alpha)$  or  $A$  be the subset  $\{x^{-i}rx^i \mid r \in R, i \geq 0\}$  of the skew Laurent polynomial ring  $R[x, x^{-1}; \alpha]$ , where  $\alpha : R \rightarrow R$  is an injective ring endomorphism of a ring  $R$ . Elements of  $R[x, x^{-1}; \alpha]$  are finite sums of elements of the form  $x^{-j}rx^i$  where  $r \in R$  and  $i, j$  are non-negative integers. Multiplication is subject to  $xr = \alpha(r)x$  and  $rx^{-1} = x^{-1}\alpha(r)$  for all  $r \in R$ . Note that for each  $j \geq 0$ ,  $x^{-i}rx^i = x^{-(i+j)}\alpha^j(r)x^{(i+j)}$ . It follows that the set  $A(R, \alpha)$  of all such elements forms a subring of  $R[x, x^{-1}; \alpha]$  with

$$\begin{aligned} x^{-i}rx^i + x^{-j}sx^j &= x^{-(i+j)}(\alpha^j(r) + \alpha^i(s))x^{(i+j)} \\ (x^{-i}rx^i)(x^{-j}sx^j) &= x^{-(i+j)}(\alpha^j(r)\alpha^i(s))x^{(i+j)} \end{aligned}$$

for  $r, s \in R$  and  $i, j \geq 0$ .

**Proposition 2.13.** *The following are equivalent for a ring  $R$ :*

- (1)  $R$  is semicommutative.
- (2)  $A(R, \alpha)$  is semicommutative.

*Proof.* (1)  $\Rightarrow$  (2). Let  $(x^{-i}rx^i)(x^{-j}sx^j) \in A(R, \alpha)$ . Suppose that  $(x^{-i}rx^i)(x^{-j}sx^j) = 0$ . Then  $x^{-(i+j)}(\alpha^j(r)\alpha^i(s))x^{(i+j)} = 0$  and so  $\alpha^j(r)\alpha^i(s) = 0$ . Hence  $\alpha^k(\alpha^j(r)\alpha^i(s)) = \alpha^{k+j}(r)\alpha^{k+i}(s) = 0$ , and  $\alpha^{j+k}(r)\alpha^{j+i}(t)\alpha^{i+k}(s) = 0$  by (1). For any  $x^{-k}tx^k \in A(R, \alpha)$

$$\begin{aligned} (x^{-i}rx^i)(x^{-k}tx^k)(x^{-j}sx^j) &= x^{-(i+k)}(\alpha^k(r)\alpha^i(t))x^{(i+k)}(x^{-j}sx^j) \\ &= x^{-(i+k+j)}\alpha^j(\alpha^k(r)\alpha^i(t))\alpha^{i+k}(s)x^{(i+k+j)} \\ &= x^{-(i+k+j)}\alpha^{j+k}(r)\alpha^{j+i}(t)\alpha^{i+k}(s)x^{(i+k+j)}. \end{aligned}$$

(2)  $\Rightarrow$  (1) From the fact that  $R \leq A(R, \alpha)$ ,  $R$  is semicommutative.  $\square$

**Proposition 2.14.** *Assume that the ring  $S = R[x]/(x^n)$  is a semicommutative ring for any  $n = 2, 3, \dots$ . Then  $R$  is a semicommutative ring.*

*Proof.* Let  $ab = 0$ . Take  $f(x) = a + (x^n), f(x) = b + (x^n) \in R[x]/(x^n) = S$ . Then

$f(x)g(x) = 0_S$ . By assumption  $(a + (x^n))(r + (x^n))(b + (x^n)) = 0_S$ . So  $arb \in (x^n)$  implies  $arb = 0$  for any  $r \in R$ .  $\square$

For a reduced ring  $R$ , it is interesting to find which subrings of  $T_n(R)$  are semicommutative. For this purpose, we introduce some notation. For number  $n \geq 4$  and any  $m$  from set  $\{1, \dots, n\}$ , we let

$$T_n^m(R) = \left\{ \sum_{i=j}^n \sum_{j=1}^m a_j E_{(i-j+1)i} + \sum_{i=j}^{n-m} \sum_{j=1}^{n-m} r_{ij} E_{j(m+i)} : a_j, r_{ij} \in R \right\},$$

where  $\{E_{i,j} : 1 \leq i, j \leq n\}$  are the matrix units. Then each element of  $T_n^m(R)$  has the matrix form

$$\begin{bmatrix} a_1 & a_2 & \dots & a_m & a_{1(m+1)} & \dots & a_{1n} \\ 0 & a_1 & \dots & a_{m-1} & a_m & \dots & a_{2n} \\ 0 & 0 & a_1 & \dots & & & a_{3n} \\ & & & \dots & & & \\ & & & & & & a_1 \end{bmatrix},$$

where  $a_1, \dots, a_m, a_{1(m+1)}, \dots, a_{(n-m)n} \in R$ .

**Theorem 2.15.** *Let  $R$  be a reduced ring. Then for number  $n \geq 4$  and  $k = [n/2]$ ,  $T_n^k(R)$  is semicommutative ring but  $T_n^{k-1}(R)$  is not.*

*Proof.* Let  $A = \sum_{i=1}^{n-k} E_{i(i+k-1)}$ ,  $B = E_{(n-k+1)n} \in T_n^{k-1}(R)$ . Then  $AB = 0$ . But for  $C = \sum_{j=i}^n \sum_{i=1}^n E_{ij} \in T_n^{k-1}(R)$ ,  $ACB \neq 0$ . So  $T_n^{k-1}(R)$  is not semicommutative. To complete the proof that  $T_n^k(R)$  is semicommutative ring for  $n \geq 4$  and  $k = [n/2]$ , it is enough to consider the case  $n = 5$ . The same proof will work for any  $n \geq 4$  and  $k = [n/2]$ . Let  $n = 5$ . Then  $k = 2$ . Let

$$A = \begin{bmatrix} a_1 & a_2 & a_{13} & a_{14} & a_{15} \\ 0 & a_1 & a_2 & a_{24} & a_{25} \\ 0 & 0 & a_1 & a_2 & a_{35} \\ 0 & 0 & 0 & a_1 & a_2 \\ 0 & 0 & 0 & 0 & a_1 \end{bmatrix}, \quad B = \begin{bmatrix} b_1 & b_2 & b_{13} & b_{14} & b_{15} \\ 0 & b_1 & b_2 & b_{24} & b_{25} \\ 0 & 0 & b_1 & b_2 & b_{35} \\ 0 & 0 & 0 & b_1 & b_2 \\ 0 & 0 & 0 & 0 & b_1 \end{bmatrix}$$

be elements of  $T_5^2(R)$  and  $AB = 0$ . We show that each term in the following system of equations obtained from  $AB = 0$  is zero:

$$A_1 B^1 : a_1 b_1 = 0 \quad \dots \quad (1)$$

$$A_1 B^2 : a_1 b_2 + a_2 b_1 = 0 \quad \dots \quad (2)$$

$$A_1 B^3 : a_1 b_{13} + a_2 b_2 + a_{13} b_1 = 0 \quad \dots \quad (3)$$

$$A_1 B^4 : a_1 b_{14} + a_2 b_{24} + a_{13} b_2 + a_{14} b_1 = 0 \quad \dots \quad (4)$$

$$A_1 B^5 : a_1 b_{15} + a_2 b_{25} + a_{13} b_{35} + a_{14} b_2 + a_{15} b_1 = 0 \quad \dots \quad (5)$$

$$A_2 B^4 : a_1 b_{24} + a_2 b_2 + a_{24} b_1 = 0 \quad \dots \quad (6)$$

$$A_2B^5 : a_1b_{25} + a_2b_{35} + a_{24}b_2 + a_{25}b_1 = 0 \quad \dots\dots\dots (7)$$

$$A_3B^5 : a_1b_{35} + a_2b_2 + a_{35}b_1 = 0 \quad \dots\dots\dots (8)$$

To prove each term in these equations( $n = 5$  or any other  $n$ ) is zero, we will proceed as follows: For  $1 \leq j \leq k$  we show all terms of  $A_1B^j$  are zero. Next for  $0 \leq i \leq n - k - 2$  we prove each term in the equations  $A_{n-k-i}B^{n-i}, A_{n-k-i}B^{n-i+1}, \dots, A_{n-k-i}B^n$  is zero. By using preceding results finally we show each term of the equations  $A_1B^{k+1}, \dots, A_1B^n$  is zero.

Note that by hypothesis from  $rs = 0$  for any  $r$  and  $s$  in  $R$  we get  $sr = 0$  and  $rRs = 0$ . Also from  $r^2s = 0$  we have  $(rs)^2 = 0$  and so  $Rs = sr = 0$ . We make use these implications without referring to the hypothesis. Now multiply (2) from left by  $a_1$ , we have  $a_1^2b_2 + a_1a_2b_1 = 0$ . By (1) and hypothesis,  $a_1a_2b_1 = 0$ . So  $a_1^2b_2 = 0$  and then  $a_1b_2 = 0$ . From (2),  $a_2b_1 = 0$ . Left multiplying (3) by  $a_1$ , we have  $a_1^2b_{13} = 0$ . Hence  $a_1b_{13} = 0$ . Then (3) becomes  $a_2b_2 + a_3b_1 = 0$ . Left multiplying this equation by  $a_2$ , we have  $a_2^2b_2 = 0 = a_2b_2$ . Hence  $a_{13}b_1 = 0$  from (3). Hence each term in the equations (1), (2) and (3) are zero.

Now we left multiply (8) by  $a_1$  and obtain  $a_1^2b_{35} = 0$  since  $a_1b_2 = 0$  and  $a_1b_1 = 0$  imply  $a_1a_2b_2 + a_1a_3b_1 = 0$ . From (8)  $a_2b_2 + a_{35}b_1 = 0$ . Left multiply the latter by  $a_2$  and use  $a_2b_1 = 0$  we get  $a_2^2b_2 = 0$ . Hence  $a_2b_2 = 0$ . By (8)  $a_{35}b_1 = 0$ .

Left multiply (6) by  $a_1$  and use  $a_1b_2 = 0$  and  $a_1b_1 = 0$  to obtain  $a_1b_{24} = 0$ . From (6)  $a_{24}b_1 = 0$  since  $a_2b_2 = 0$ .

Left multiply (7) by  $a_1$  and use  $a_1b_{35} = 0$ ,  $a_1b_2 = 0$  and  $a_1b_1 = 0$  to obtain  $a_1b_{25} = 0$ . (7) induces to  $a_2b_{35} + a_{24}b_2 + a_{25}b_1 = 0$ . We left multiply the latter by  $a_2$  to obtain  $a_2b_{35} = 0$ . From (7) we have  $a_{24}b_2 + a_{25}b_1 = 0$ . Left multiply this by  $b_1$  and use  $a_{24}b_1 = 0$  to obtain  $b_1^2a_{25} = 0$ . Hence  $b_1a_{25} = 0$ . Now we go to the equation (5) to left multiply it by  $a_1$  and use  $a_1b_{35} = 0$ ,  $a_1b_{25} = 0$ ,  $a_1b_2 = 0$ ,  $a_1b_1 = 0$  to get  $a_1^2b_{15} = 0$ . Hence  $a_1b_{15} = 0$ . From (5) we have  $a_2b_{25} + a_{13}b_{35} + a_{14}b_2 + a_{15}b_1 = 0$ . Similarly this procedure continues to obtain each term in the latter equation is zero:  $a_2b_{25} = 0$ ,  $a_{13}b_{35} = 0$ ,  $a_{14}b_2 = 0$ ,  $a_{15}b_1 = 0$ . As for (4), left multiply it by  $a_1$  to get  $a_1^2b_{14} = 0$  since  $a_1b_{24} = 0$ ,  $a_1b_2 = 0$  and  $a_1b_1 = 0$ . So  $a_1b_{14} = 0$ . From (4)  $a_2b_{24} + a_{13}b_2 + a_{14}b_1 = 0$ . Left multiply it by  $a_2$  and use  $a_2b_2 = 0$  and  $a_2b_1 = 0$  to obtain similarly  $a_2b_{24} = 0$ . We are left with  $a_{13}b_2 + a_{14}b_1 = 0$ . Left multiply it by  $b_2$  and use  $a_{14}b_2 = 0$  to obtain  $b_2^2a_{13} = 0$ . Hence  $b_2a_{13} = 0$ . Thus  $a_{14}b_1 = 0$ . Since  $R$  is semicommutative, the rest of the proof is clear.  $\square$

**Corollary 2.16.** *Let  $R$  be a prime ring. Then  $R[x]/(x^n)$  is Armendariz if and only if  $R[x]/(x^n)$  is semicommutative.*

*Proof.* Clear from [9, Corollary 1.5] and Theorem 2.23.  $\square$

**Corollary 2.17.** *If  $R$  is a reduced ring then  $R[x]/(x^n)$  is semicommutative.*

**Corollary 2.18.** *Let  $R$  be an Armendariz ring. Then  $R$  is semicommutative if and only if  $R[x]/(x^n)$  is semicommutative.*

**Acknowledgement.** It is a pleasure to thank Professor S. Tariq Rizvi for his helpful suggestions.

## References

- [1] M. Baser and N. Agayev, *On Reduced and Semicommutative Modules*, Turk. J. Math., **30**(2006), 285-291.
- [2] G. F. Birkenmeier, J. Y. Kim and J. K. Park, *Principally Quasi-Baer Rings*, Comm. Algebra, **29**(2001), 639-660.
- [3] A. M. Buhphang and M. B. Rege, *Semicommutative Modules and Armendariz Modules*, Arab Journal of Mathematical Sciences, **8**(June 2002), 53-65.
- [4] C. Huh, Y. Lee and A. Smoktunowicz, *Armendariz Rings and Semicommutative Rings*, Comm. Algebra, **30**(2002), 751-761.
- [5] D. A. Jordan, *Bijjective Extension of Injective Ring Endomorphisms*, J. London Math. Soc., **35**(2)(1982), 435-448.
- [6] N. K. Kim and Y. Lee, *Armendariz Rings and Reduced Rings*, J. Algebra, **223**(2000), 477-488.
- [7] I. Kaplansky, *Rings of operators*, W. A. Benjamin, New York, 1968.
- [8] T. K. Lee and Y. Zhou, *Reduced Modules, Rings, modules, algebras and abelian groups*, 365-377, Lecture Notes in Pure and Appl. Math., 236, Dekker, New york, (2004).
- [9] T. K. Lee and Y. Zhou, *Armendariz and Reduced Rings*, Comm. Algebra, **32**(6)(2004), 2287-2299.
- [10] M. B. Rege and S. Chhawchharia, *Armendariz rings*, Proc. Japan Acad., **73**(A)(1997), 14-17.
- [11] S. T. Rizvi and C. S. Roman, *Baer and Quasi-Baer Modules*, Comm. Algebra, **32**(2004), 103-123.