

Comparison of the classification ratios of four depression rating scales commonly used in Turkey

Gaye Saltukoğlu & Arkun Tatar

To cite this article: Gaye Saltukoğlu & Arkun Tatar (2017) Comparison of the classification ratios of four depression rating scales commonly used in Turkey, *Psychiatry and Clinical Psychopharmacology*, 27:1, 76-83, DOI: [10.1080/24750573.2017.1293250](https://doi.org/10.1080/24750573.2017.1293250)

To link to this article: <https://doi.org/10.1080/24750573.2017.1293250>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 31 Mar 2017.



Submit your article to this journal [↗](#)



Article views: 899



View related articles [↗](#)



View Crossmark data [↗](#)

Comparison of the classification ratios of four depression rating scales commonly used in Turkey*

Gaye Saltukoğlu^a and Arkun Tatar^b

^aDepartment of Psychology, Fatih Sultan Mehmet Vakif University, Istanbul, Turkey; ^bDepartment of Psychology, The University of Texas, Austin, TX, USA

ABSTRACT

Objective: According to literature more than 20 depression scales are in use in Turkey. Considering that depression is a popular area of study, it may not seem abnormally unusual that there are so many measuring scales available. However, so many measuring instruments may lead to a problem of different sensitivity levels and raise the question of whether or not all the instruments have the same sensitivity in measuring the particular entity. The purpose of this study is to compare the four commonly used self-report scales adapted into Turkish, namely CES-Depression Scale, Beck Depression Inventory, the Zung Self-Rating Depression Scale, and the Hospital Anxiety and Depression Scale (depression subscale) by cross-validation. **Method:** These depression scales had been applied to three hundred and forty-one subjects and total scores of the subjects for each scale have been obtained. Next, the sample group was divided into two according to group averages of total scale scores. Normative scores and cut-off scores have not been considered because the study objective was to compare these scales on a theoretical basis. The groups below and above average for each of the four scales have been compared by the ROC curve analyzes.

Results: The results showed that the total score of Beck Depression Inventory had been grouped correctly by the Zung Self-Rating Depression Scale at a ratio of 0.871, the Hospital Anxiety and Depression Scale (depression subscale) at a ratio of 0.885, and by CES-Depression Scale at a ratio of 0.874. The total score of CES-Depression Scale had been correctly grouped by Beck Depression Inventory at a ratio of 0.871, the Zung Self-Rating Depression Scale at a ratio of 0.869, and by the Hospital Anxiety and Depression Scale (depression subscale) at a ratio of 0.862. The total score of the Zung Self-Rating Depression Scale has been correctly grouped by the Hospital Anxiety and Depression Scale depression subscale at a ratio of 0.848, Beck Depression Inventory at a ratio of 0.872, and by CES-Depression Scale at a ratio of 0.878. The total score of the Hospital Anxiety and Depression Scale (depression subscale) has been correctly grouped by the Zung Self-Rating Depression Scale at a ratio of 0.848, Beck Depression Inventory at a ratio of 0.889, and by CES-Depression Scale at a ratio of 0.887.

Conclusion: The overall results showed that the scales cross-validated with ratios ranging from 0.85 to 0.89. The classifying ratios obtained by ROC analysis were similar across four depression scales.

ARTICLE HISTORY

Received 15 October 2015
Accepted 5 January 2016

KEYWORDS



Cross-validity; ROC curve; depression scale

Introduction

Symptoms of depression have been described roughly, and include a wide range of manifestations, ranging from short-term distress to despair, guilt, unwillingness, and low self-esteem. Long-term depression causes fatigue, sleeplessness, chronic pain, and extreme weight loss or gain [1,2]. Depression has been related to socioeconomic status, ethnicity, age, and gender in various studies, and females, lower socioeconomic groups, and people in the age range of 25–34 are the most affected ones [3–7]. The prevalence of depression is reported to be 15% in women and 10% in men [8]. Moreover, this disorder is related to one's livelihood

[9]; thus it is estimated that by 2030 depression would be the most important reason for labor loss [4,10]. Depression is the most common and widespread psychological disorder and occurs in almost every society [4]; however, in underdeveloped and developing countries, it is difficult to determine the prevalence due to the lack of adapted scales and unestablished reliability and validity [11].

Tests labeled “Depression Tests” are claimed to measure similar constructs and are designed to provide a total individual score [12]. Santor et al. [13] reported that between the years of 1918 and 2006 more than 280 depression scales have been developed and published,

CONTACT Arkun Tatar  arkuntatar@yahoo.com  Department of Psychology, The University of Texas at Austin, 108 E. Dean, Keeton Stop A8000 SEA 3.402 Austin, TX 78712-1043, USA

*An early version of this paper was first presented at the International Psychological Applications Conference and Trends (InPACT) in Ljubljana, Slovenia, May 2015.

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

with an increased rate of development after 1940. Scales have been developed on different samples and in different contexts, which have been categorized based on different theoretical backgrounds, such as psychoanalytic, cognitive, behavioral, and evolutionary as well as those related to attachment and interpersonal relationships. Most of these tests have been developed using classical test theory and focus on a “real” score and measurement error.

Standard depression tests are known to be useful when used for screening purposes in mental health research and in general populations. It is also stated that such measuring instruments possess many favorable characteristics, such as good psychometric qualities, convenient number of items, being understandable, free of charge, and easily accessible [14,15].

In an earlier study, Ceyhun [16] reported that 13 depression scales have been used in Turkey, 2 of which are applied by clinicians, 9 of them consist of self-report questionnaires, and 2 of them are structured interview scales.

In addition, the CES-Depression Scale [17], the Calgary Depression Scale [18], the Cornell Scale for Depression in Dementia [19,20], Edinburgh Postnatal Depression Scale [21], Geriatric Depression Scale [22], Postpartum Depression Screening Scale [23], the Boratav Depression Screening Scale [24], and General Health Questionnaire-12 [25] are some of the scales which have been translated and adapted into Turkish.

In studies where Turkish versions of depression scales are used, depression levels have usually been examined on special groups [26] or the test itself has been validated [27] or depression has been compared with some other theoretical construct/concept [28,29]. There are very few studies other than adaptation of the scales into Turkish or examination of psychometric properties [30–32]. Likewise, there are few studies where the results of different depression scales are head-to-head compared [33–40].

Considering that depression is a serious medical illness with important societal implications, it may not seem unusual that there are many measuring scales available. However, with so many instruments available, it is important to determine if the instruments provide comparable measurement of the construct and have similar sensitivity. In addition, depression total test scores developed by classical test theory cannot be compared with others in terms of test equivalence [12]. Therefore, the criteria for selecting and using one instrument over another in a sample from Turkey are currently unclear. Given these factors, the purpose of this study was to compare the Turkish version of four self-report depression scales, namely Beck Depression Inventory (BDI), the Zung Self-Rating Depression Scale (SDS), the Hospital Anxiety and

Depression Scale (HADS), and the CES-Depression Scale (CES-D) by a cross-validation study using Receiver Operating Characteristics (ROC) curves. In this way, we aimed to obtain evidence to determine if these instruments measure the same construct and whether one can be substituted for another.

Methods

Participants

The sample included 235 women (68.9%) and 106 men (31.1%) ranging in age from 18 to 62 (mean = 29.26 ± 10.79 years). Participants consisted of 32 elementary school graduates (9.4%), 19 middle school graduates (5.6%), 73 high school graduates (21.4%), and 211 university graduates (61.9%) (6 people did not specify their level of education (1.8%)). Of all the participants, 60.1% ($n = 205$) were single, 34.0% ($n = 116$) married, 1.5% ($n = 5$) divorced, and 2.6% ($n = 9$) widowed (6; 1.8% of participants did not specify their marital status). Finally, economic status of the participants was as follows: 7.3% ($n = 25$) low income, 60.7% ($n = 207$) medium income, and 29.0% ($n = 99$) high income (10; 2.9% of participants did not specify their income).

Instruments

The CES-Depression Scale

The scale, which has been widely used and translated into several languages [41] was developed by Radloff [42] to examine the items of available depression scales, with the purpose of screening depressive symptoms in the general population. The scale assesses emotional and somatic symptoms of the participants in the previous week [42]. The scale has been translated into Turkish by Tatar and Saltukoğlu [17]. The scale consists of 20 items scored between 0 and 3 and the score obtained varies between 0 and 60. Although there is not a defined factor construct of the scale, usually Radloff's proposed four-factor construct [42] has been supported [3,5,43]. However, researchers have reported different numbers of factor constructs [44]. The scale has been one of the most widely used scales in studies related to the treatment and prognosis of depression [45].

The Beck Depression Inventory

This inventory was developed in 1961 by Beck et al. [46] and consists of 21 items scored between 0 and 3 on a four-point rating scale and the obtained scores vary between 0 and 63. Items of the inventory focus on emotional, behavioral, and somatic symptoms [5]. This test is the most widely used scale for the measurement of depression [1]. The scale was adapted into Turkish by two independent studies [16]. In the

present study, the form that has been used is the one adapted into Turkish by Hisli [47–49].

In some of the studies using the Turkish form of the scale topics such as major depression [50], postpartum depression [36], somatic symptoms [38], and comorbidity of impulse control disorder in depressive patients [35], depression in coronary artery disease [26], essential tremor, Parkinson's disease [51], and in haemodialysis patients [52], relationship between pain and depression in patients with knee osteoarthritis [53], depression in adolescents three and a half years after the Marmara earthquake [54], depression in mothers of children with food refusal [55], dimensions of alexithymia and the intensity of depression [56], relationship between depression and anger in patients with antisocial personality disorder [57] have been examined.

The Zung Self-Rating Depression Scale

This scale, developed by Zung [58], consists of 20 items scored between 1 and 4 on a four-point rating scale and obtained total score varies between 20 and 80. The scale was adapted into Turkish by Ceyhun and Akça [59] and measures common clinical symptoms of which half of the expressions are negative and half of them positive. The scale measures psychological and somatic symptoms of depression. Although many of the studies have not specified defined sub-dimensions or factor constructs, the scale has been widely defined by two factors; well-being and depressive symptoms (or positive and negative symptoms) [5].

In some of the studies where Beck Depression Inventory and the Hospital Anxiety and Depression Scale are used, the Turkish form of this scale of depression levels in mothers of handicapped children [40] and low back pain and its relationship with pain-related disability and depression have been examined [60].

The Hospital Anxiety and Depression Scale (depression subscale)

This scale, developed by Zigmond and Snaith [61], was adapted into Turkish by Aydemir et al. [62]. The scale consists of a total of 14 items of which 7 items are for the assessment of anxiety (The Hospital Anxiety and Depression Scale-A; odd numbered items), and the remaining 7 items are for the assessment of depression (The Hospital Anxiety and Depression Scale-D; even numbered items). Both of the subscales are scored between 0 and 3 on a 4-point scale and the total score obtained ranges between 0 and 21 [62]. In this study, 14 items were administered to all participants to maintain the integrity of the test; however, only the items of the depression sub-dimension have been assessed.

Some of the other studies using the Turkish version of the Hospital Anxiety and Depression Scale examined

the relationship between depression and alopecia areata and alexithymia [63], examined depression in women in the premenopausal and postmenopausal period [64] and nurses [65], in patients with diabetes mellitus [66,67], in patients with low back and neck pain [68], and in medical inpatients [69].

Procedure

Subjects were selected by convenience sampling and participated in the study voluntarily. Subjects were asked whether they wanted to participate in a scientific survey and they were informed that they can withdraw their participation at any time. One participant did not complete both the CES-Depression Scale and the Zung Self-Rating Depression Scale, one participant did not complete the Beck Depression Inventory, one participant did not complete the Zung Self-Rating Depression Scale, and one participant did not complete the Hospital Anxiety and Depression Scale. Nonetheless, these participants completed and returned all other scales. All scales were administered on an individual basis. It took 10–15 minutes for a participant to complete all the forms.

Statistical analysis

First, the total scores for each scale were calculated. From the total scores, the mean scores for each scale were determined and are shown in Table 1. Next, the sample group is divided into two according to group averages of total scale scores. Normative scores and cutting points have not been considered because the purpose here is to compare the scales on a theoretical basis. The groups below and above average for each four of the scales have been compared by ROC curves.

Results

Cronbach's alpha internal consistency coefficients were calculated and were shown in the same table. Results showed that Cronbach's alpha internal consistency coefficients are 0.90 for the CES-Depression Scale, 0.91 for the Beck Depression Inventory, 0.84 for the Zung Self-Rating Depression Scale, and 0.85 for the Hospital Anxiety and Depression Scale-D.

The study group was then divided into two groups, representing participants who scored above and below the group mean. For this procedure, the norms and the cut-off points of the scales were not taken into account because the purpose was to compare the scales on a theoretical basis. Table 2 shows the comparisons of the groups that were derived from each scale, where one can see the curve ratios of people who are above and below the mean with respect to their grouping in each of the other scales. Out of the people who are situated below the mean in the CES-Depression Scale test

Table 1. Mean values and reliability coefficients of the scales.

	<i>n</i>	No. of items	Cronbach's alpha	Minimum	Maximum	Mean	Standard deviation
CES-Depression Scale	340	20	0.90	0	60	17.34	12.18
Beck Depression Inventory	340	21	0.91	0	63	10.29	9.23
The Zung Self-Rating Depression Scale	339	20	0.84	20	76	39.65	10.90
The Hospital Anxiety and Depression Scale-D	340	7	0.85	0	21	5.81	4.64

($n = 189$), 32 of them were above the mean in the Beck Depression Inventory, 45 of them were above the mean in the Zung Self-Rating Depression Scale, and 42 of them were above the mean in the Hospital Anxiety and Depression Scale-D. The distribution of these values has been compared by ROC curves. It is sufficient to give the number and percentage distributions of the scales relative to each other.

Next, Pearson's correlation coefficients of the scales with respect to each other were calculated and are shown in Table 3. The correlation coefficients found between the scales ranged between 0.75 and 0.80. The Zung Self-Rating Depression Scale obtained both the lowest and the highest scores, where it showed the highest correlation with the CES-Depression Scale and the lowest correlation with the Beck Depression Inventory.

Finally, scores of each scale below and above the group mean scores with respect to total scale scores were compared with the total scores of the other scales by using the ROC curve.

In Figure 1 results of the ROC curve showed that the CES-Depression Scale correctly classifies the Beck Depression Inventory by a ratio of 0.871, the Zung Self-Rating Depression Scale by a ratio of 0.869, and the Hospital Anxiety and Depression Scale-D by a ratio of 0.862.

In Figure 2, results of the ROC curve showed that the Beck Depression Inventory correctly classified the CES-Depression Scale by a ratio of 0.874, the Zung

Self-Rating Depression Scale by a ratio of 0.871, and the Hospital Anxiety and Depression Scale-D by a ratio of 0.885.

In Figure 3, results of the ROC curve show that the Zung Self-Rating Depression Scale correctly classifies the CES-Depression Scale by a ratio of 0.878, the Beck Depression Inventory by a ratio of 0.872, and the Hospital Anxiety and Depression Scale-D by a ratio of 0.848.

In Figure 4, results showed that the Hospital Anxiety and Depression Scale-D correctly classified the CES-Depression Scale by a ratio of 0.887, the Beck Depression Inventory by a ratio of 0.889, and the Zung Self-Rating Depression Scale by a ratio of 0.848.

Our results showed that groups located below and above the group mean scores had substantially similar ratios for correctly classifying the other three scales according to the ROC analysis. For the CES-Depression Scale, the average correct classifying ratio of other three scales is 0.867; for the Beck Depression Inventory, it is 0.877; for the Zung Self-Rating Depression Scale; it is 0.866; and for The Hospital Anxiety and Depression Scale-D, it is 0.872.

Discussion

Many scales have been developed for the purpose of measuring and assessing depression. In this study, only four of the most commonly used scales in Turkey

Table 2. The comparison of groups with below and above means in four scales.

		<i>n</i>	Beck Depression Inventory		The Zung Self-Rating Depression Scale		The Hospital Anxiety and Depression Scale-D		Total
			BA	AA	BA	AA	BA	AA	
			%	%	%	%	%	%	
CES-Depression Scale	BA	<i>n</i>	157	32	144	45	147	42	189
		%	83.1	16.9	76.2	23.8	77.8	22.2	100.0
	AA	<i>n</i>	47	103	25	125	31	119	150
		%	31.3	68.7	16.7	83.3	20.7	79.3	100.0
Total	<i>n</i>	204	135	169	170	178	161	339	
	%		60.2	39.8	49.9	50.1	52.5	47.5	100.0
Beck Depression Inventory	BA	<i>n</i>			148	56	157	48	205
		%			72.5	27.5	76.6	23.4	100.0
	AA	<i>n</i>			21	113	22	112	134
		%			15.7	84.3	16.4	83.6	100.0
Total	<i>n</i>			169	169	179	160	339	
	%			50.0	50.0	52.8	47.2	100.0	
The Zung Self-Rating Depression Scale	BA	<i>n</i>					133	36	169
		%					78.7	21.3	100.0
	AA	<i>n</i>					44	125	169
		%					26.0	74.0	100.0
Total	<i>n</i>					177	161	338	
	%					52.4	47.6	100.0	

Note: BA: below-average group; AA: above-average group.



Table 3. Correlation coefficients and inter-correlations of depression scales.

		Beck Depression Inventory	The Zung Self-Rating Depression Scale	The Hospital Anxiety and Depression Scale-D
CES-Depression Scale	<i>r</i>	0.79***	0.80***	0.77***
	<i>n</i>	339	339	339
Beck Depression Inventory	<i>r</i>		0.75***	0.78***
	<i>n</i>		338	339
The Zung Self-Rating Depression Scale	<i>r</i>			0.76***
	<i>n</i>			338


*** $p < .001$.

were compared. Without considering norms and cut-off scores, classification ratios were roughly compared. In our sample, participants completed these depression inventories and the results indicated that the four scales showed similar Cronbach's alpha coefficients ranging between 0.84 and 0.91. Likewise, the total scores of the scales were similar, yielding correlation coefficients that ranged between 0.75 and 0.80.

When we applied the ROC analysis to test whether the groups that are divided below and above the total score averages correctly classified the other scales, we found that the CES-Depression Scale classified other scales with the mean value of 0.867, the Beck Depression Inventory by 0.877, the Zung Self-Rating Depression Scale by 0.866, and the Hospital Anxiety and Depression Scale-D by 0.872. These values are very close to each other. Although the Zung Self-Rating Depression Scale's correct classification ratio of the Hospital Anxiety and Depression Scale-D and the Hospital Anxiety and Depression Scale-D's correct classification ratio were smaller than the others by a margin, the differences were small when the average values are considered.

**Figure 1.** Comparison of the CES-Depression Scale with other scales' total scores by ROC curve.**Figure 2.** Comparison of the Beck Depression Inventory with other scales' total scores by ROC curve.

ROC analysis is a statistical technique that shows the sensitivity and specificity of the test graphically. This analysis has been reported as an effective technique to evaluate the performance of diagnostic tests [70,71]. In this present study, we evaluated the depression scales using ROC analyzes. However, ROC analysis in this study was not used to evaluate the diagnostic power of the scales in depressed patients, but to compare the classifying ratios of the scales with each other. The results obtained for each scale showed at what ratio a scale reflected the other three scales. In other words, the result obtained for each test is the ratio of overlap with the other three scales in terms of diagnosis. Results showed that the differences between the

**Figure 3.** Comparison of the Zung Self-Rating Depression Scale with other scales' total scores by ROC curve.




Figure 4. Comparison of the Hospital Anxiety and Depression Scale-D with other scales' total scores by ROC curve.

four depression scales were extremely small. This result explains that the use of these scales is interchangeable. On the other hand, whether a depression scale determines the level of depression in a person or makes a distinction between depressive and non-depressive individuals is beyond the scope of this study.

Due to different number of items and different ways of scoring, the total scores of these four scales are not isomorphic. That is, 10 points obtained from the Hospital Anxiety and Depression Scale-D and 10 points obtained from CES-D do not show the same depression level. While the Hospital Anxiety and Depression Scale-D with 7 items gives a total score between 0 and 21, the CES-Depression Scale with 20 items gives a total score between 0 and 60. For this reason, the average scores of each scales which were obtained on the same group were used as cut-off scores. Even if the scale scores are mathematically isomorphic, in terms of measured variance of scale scores, that is, in terms of depression levels, these test scores will not be enough for ensuring measurement equivalence. There are some psychometric techniques to determine measurement equivalence. However, in this study, the way of the use of ROC analysis to compare tests in a sense resolves the problem without resorting to such psychometric techniques. In other words, the technique discussed in this study has made it possible for the tests to be compared to each other without ensuring measurement equivalence of total scores of the tests.

This study has certain limitations. Although the study group is small, it is possible to conclude that these scales do not produce different classifications and can be used interchangeably in screening studies. Although the focal point of this study was not the length or the number of test items, it is worth pointing

out that only the Hospital Anxiety and Depression Scale-D subscale was used in this study, which consisted of only seven items and was significantly shorter than the others. Notwithstanding, considering that correct classifying ratio of this subscale was not significantly different from the other three tests, it can be concluded that the Hospital Anxiety and Depression Scale-D subscale is useful for screening purposes in studies with large samples apart from the purpose of individual diagnosis.

Another important limitation of this study is that no comparisons were made between clinical and non-clinical samples. The depression scales used in this study are also used as clinical diagnostic tools besides their use for research purposes. For this reason, the examination of these scales on the participants that represent both clinical and non-clinical groups would be useful. We have previously indicated that the criteria by which depression scales are selected in Turkey are not clear. We, therefore, selected four tests based on their widespread use in Turkey. We are unable to provide clear selection criteria for the scales under study. Despite these limitations, each of the scales under investigation in this study successfully classified the other three tests when below and above average scale scores were compared. The classifying ratios obtained by ROC analysis were similar across four depression scales. We conclude that these tests can be used interchangeably in Turkey.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

- [1] Spielberger CD, Ritterband LM, Reheiser EC, et al. The nature and measurement of depression. *Int J Clin Health Psychol.* 2003;3(2):209–234.
- [2] de Guzman AB, Lagdaan LFM, Lagoy MLV. The role of life-space, social activity, and depression on the subjective memory complaints of community-dwelling Filipino elderly: a structural equation model. *Educ Gerontol.* 2014;41(5):1–13.
- [3] Kim G, Decoster J, Huang CH, et al. Race/ethnicity and the factor structure of the center for epidemiologic studies depression scale: a meta-analysis. *Cultur Divers Ethnic Minor Psychol.* 2011;17(4):381–396.
- [4] Missinne S, Vandeviver C, Van de Velde S, et al. Measurement equivalence of the CES-D 8 depression-scale among the ageing population in eleven European countries. *Soc Sci Res.* 2014;46:38–47.
- [5] Shafer AB. Meta-analysis of the factor structures of four depression questionnaires: Beck, CES-D, Hamilton, and Zung. *J Clin Psychol.* 2006;62(1):123–146.
- [6] Sigurdsson B, Palsson SP, Aevarsson O, et al. Validity of gotland male depression scale for male depression in a community study: the Sudurnesjamenn study. *J Affect Disord.* 2015;173:81–89.
- [7] Zhong Q, Gelaye B, Rondon M, et al. Comparative performance of patient health questionnaire-9 and

- Edinburgh postnatal depression scale for screening antepartum depression. *J Affect Disord.* 2014;162:1–7.
- [8] Regier DA, Boyd JH, Burke JD Jr, et al. One-month prevalence of mental disorders in the U.S. based on five epidemiological catchment area sites. *Arch Gen Psychiatry* 1988;45:977–986.
- [9] Fukukawa Y, Nakashima C, Tsuboi S, et al. The impact of health problems on depression and activities in middle-aged and older adults: age and social interactions as moderators. *J Gerontol B Psychol Sci Soc Sci.* 2004;59(1):19–26.
- [10] Chavez CA, Ski CF, Thompson DR. Psychometric properties of the cardiac depression scale: a systematic review. *Heart Lung Circ.* 2014;23(7):610–618.
- [11] Guerra M, Ferri C, Llibre J, et al. Psychometric properties of Euro-D, a geriatric depression scale: a cross-cultural validation study. *BMC Psychiatry.* 2015;15(1):1–14.
- [12] Fischer HF, Tritt K, Klapp BF, et al. How to compare scores from different depression scales: equating the patient health questionnaire (PHQ) and the ICD 10 symptom rating (ISR) using item response theory. *Int J Methods Psychiatr Res.* 2011;20(4):203–214.
- [13] Santor DA, Gregus M, Welch A. Focus article: eight decades of measurement in depression. *Measurement (Mahwah NJ).* 2006;4(3):135–155.
- [14] Matijasevich A, Munhoz TN, Tavares BF, et al. Validation of the Edinburgh postnatal depression scale (EPDS) for screening of major depressive episode among adults from the general population. *BMC Psychiatry.* 2014;14(284):1–9.
- [15] Stukenberg KW, Dura JR, Kiecolt-Glaser JK. Depression screening scale validation in an elderly, community-dwelling population. *Psychol Assess.* 1990;2(2):134–138.
- [16] Ceyhan B. [Psychological assessment of depression]. *Duygudurum Dizisi.* 2001;1(4):192–197. Turkish.
- [17] Tatar A, Saltukoglu G. The adaptation of the CES-depression scale into Turkish through the use of confirmatory factor analysis and item response theory and the examination of psychometric characteristics. *Klin Psikofarmakol B.* 2010;20(3):213–227.
- [18] Aydemir O, Danacı EA, Deveci A, et al. [Reliability and validity of the Turkish version of the calgary depression scale in schizophrenia]. *Arch Neuropsychiatry.* 2000;37(2):82–86. Turkish.
- [19] Amuk T, Karadag F, Oguzhanoglu N, et al. [Reliability and validity of the Cornell scale for depression in dementia in an elderly Turkish population]. *Turk Psikiyatri Derg.* 2003;14(4):263–271. Turkish.
- [20] Taycan O, Eker E. [Reliability and validity of the Cornell scale for depression in dementia in Turkish demented patients]. *Klinik Psikiyatri Dergisi.* 2009;12(1):21–34. Turkish.
- [21] Kilinc S, Torun F. [Depression rating scales used in clinical practice in Turkey]. *Dirim Tıp Gazetesi.* 2011;86(1):39–47. Turkish.
- [22] Ertan T, Eker E, Sar V. [Reliability and validity of the geriatric depression scale in an elderly Turkish population]. *Arch Neuropsychiatry.* 1997;34(2):62–71. Turkish.
- [23] Karacam Z, Kitis Y. [The postpartum depression screening scale: its reliability and validity for the Turkish population]. *Turk Psikiyatri Derg.* 2008;19(2):187–196. Turkish.
- [24] Boratav C. [The Boratav depression screening scale (Bordepta): a sensitive scale for recognizing depression in epidemiologic studies and primary health care]. *Turk Psikiyatri Derg.* 2003;14(3):172–183. Turkish.
- [25] Kilic C. [General health questionnaire: study of reliability and validity]. *Turk Psikiyatri Derg.* 1996;7(1):3–9. Turkish.
- [26] Sunbul M, Bozkurt-Zincir S, Durmus E, et al. Anxiety and depression in patients with coronary artery disease. *Klin Psikofarmakol B.* 2013;23(4):345–352.
- [27] Tatar A, Kayiran SM, Saltukoglu G, et al. Analysis of the center for epidemiologic studies depression scale (CES-D) in children and adolescents from the perspective of the item response theory. *Klin Psikofarmakol B.* 2013;23(3):242–253.
- [28] Arkar H. [The relationship between depressive mood and personality: comparison of the Beck depression inventory and the temperament and character inventory]. *New Symposium.* 2010;48(2):116–121. Turkish.
- [29] Boysan M, Besiroglu L, Kara H, et al. [Associations between the levels of depression and the sense of limit-edness: a psychometric analysis]. *Anadolu Psikiyatri Derg.* 2008;9(4):191–196. Turkish.
- [30] Karacam Z, Oner H. [Calculation of cut off points in the Turkish version of postpartum depression screening scale]. *J Anatolia Nurs Health Sci.* 2008;11(2):61–71. Turkish.
- [31] Ozer AK, Ozer S. [Factorial structure of state-trait depression scales]. *Doğuş University J.* 2011;7(2):210–217. Turkish.
- [32] Uluc S. [Examination of the criterion validity of the MMPI-2 depression, anxiety, and anger content scales]. *Turk Psikiyatri Derg.* 2008;19(1):57–66. Turkish.
- [33] Akkaya C, Eker SS, Sarandol A, et al. [The predictors of remission in major depressive disorder]. *Arch Neuropsychiatry.* 2013;50(2):122–129. Turkish.
- [34] Alkan H, Sarsan A, Topuz O, et al. [Effectiveness of home rehabilitation programme in chronic obstructive pulmonary disease]. *Turkish J Phys Med and Rehab.* 2006;52(2):51–54. Turkish.
- [35] Annagur BB, Tamam L. [Comorbidity of impulse control disorders among patients with major depression]. *Arch Neuropsychiatry.* 2011;48(1):22–30. Turkish.
- [36] Ayvaz S, Hocaoglu C, Tiryaki A, et al. [Incidence of postpartum depression in Trabzon province and risk factors at gestation]. *Turk Psikiyatri Derg.* 2006;17(4):243–251. Turkish.
- [37] Cilan H, Oymak O, Turan T, et al. [Nutritional status and depressive disorders in hemodialysis patients at Erciyes University, faculty of medicine, hospital hemodialysis unit]. *Erciyes Med J.* 2009;31(3):237–243. Turkish.
- [38] Gulec H, Sayar K, Ozkorumak E. [Somatic symptoms of depression]. *Turk Psikiyatri Derg.* 2005;16(2):90–96. Turkish.
- [39] Ozguven HD. [Prevalence of faulty automatic thoughts and dysfunctional attitudes in psychiatric crisis cases]. *Kriz Dergisi.* 1999;7(2):9–16. Turkish.
- [40] Tatar A, Kayiran SM, Saltukoglu G, et al. Examination of depression levels in mothers of handicapped children. *HealthMED.* 2011;5(1):95–100.
- [41] Boyd RC, Le H, Somberg R. Review of screening instruments for postpartum depression. *Arch Womens Ment Health.* 2005;8(3):141–153.
- [42] Radloff LS. The CES-D scale a self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977;1(3):385–401.

- [43] Verhoeven M, Sawyer MG, Spence SH. The factorial invariance of the CES-D during adolescence: are symptom profiles for depression stable across gender and time? *J Adolesc.* 2013;36(1):181–190.
- [44] Edwards MC, Cheavens JS, Heiy JE, et al. A reexamination of the factor structure of the center for epidemiologic studies depression scale: is a one-factor model plausible? *Psychol Assess.* 2010;22(3):711–715.
- [45] Aydemir O. [Therapeutic goals, clinical indicators and follow up in major depressive disorder]. *Klin Psikofarmakol B.* 2011;21(Suppl. 1):S1–S9. Turkish.
- [46] Beck AT, Ward CH, Mendelson M, et al. An inventory for measuring depression. *Arch Gen Psychiatry.* 1961;4(6):561–571.
- [47] Ulusoy M, Sahin NH, Erkmén H. Turkish version of the Beck anxiety inventory: psychometric properties. *J Cogn Psychother.* 1998;12(2):163–172.
- [48] Hisli N. [A study towards to validity of Beck depression inventory]. *Psikoloji Dergisi.* 1988;6(22):118–126. Turkish.
- [49] Hisli N. [Reliability and validity of the Beck depression inventory in university students]. *Psikoloji Dergisi.* 1989;7(23):3–13. Turkish.
- [50] Aydemir C, Temiz HV, Goka E. [Cognitive and emotional factors in major depression and suicide]. *Turk Psikiyatri Derg.* 2002;13(1):33–39. Turkish.
- [51] Yaka E, Aysevener EO, Cifcioglu O, et al. [A comparison between frequency of depressive symptoms and quality of life among patients with essential tremor and Parkinson's disease]. *Arch Neuropsychiatry.* 2011;48(4):255–260. Turkish.
- [52] Baydogan M, Dag I. [Prediction of depressiveness by locus of control, learned resourcefulness and socio-tropy-autonomy in hemodialysis patients]. *Turk Psikiyatri Derg.* 2008;19(1):19–28. Turkish.
- [53] Cetin N, Oztup P, Bayramoglu M, et al. [Relation between pain, disability and depression in patients with knee osteoarthritis]. *Turk J Rheumatol.* 2009;24(4):196–201. Turkish.
- [54] Karakaya I, Agaoglu B, Coskun A, et al. [The symptoms of PTSD, depression and anxiety in adolescent students three and a half years after the Marmara earthquake]. *Turk Psikiyatri Derg.* 2004;15(4):257–263. Turkish.
- [55] Unlu G, Aras S, Guvenir T, et al. [Family functioning, personality disorders, and depressive and anxiety symptoms in the mothers of children with food refusal]. *Turk Psikiyatri Derg.* 2006;17(1):12–21. Turkish.
- [56] Motan I, Gencoz T. [The relationship between the dimensions of alexithymia and the intensity of depression and anxiety]. *Turk Psikiyatri Derg.* 2007;18(4):333–343. Turkish.
- [57] Turkcapar H, Guriz O, Ozel A, et al. [Relationship between depression and anger in patients with antisocial personality disorder]. *Turk Psikiyatri Derg.* 2004;15(2):119–124. Turkish.
- [58] Zung WW. A self-rating depression scale. *Arch Gen Psychiatry.* 1965;12(1):63–70.
- [59] Ceyhun B, Akca F. [A study towards to validity and reliability of Zung depression scale]. In: Topsever Y, Goregenli M, editors. [Proceedings of the VIII. National Psychology Congress Scientific Studies]. 1994 Sep 21–23; Izmir. Ankara (Turkey): Turkish Psychological Association; 1996. p. 20–26. Turkish.
- [60] Tucer B, Yalcin BM, Ozturk A, et al. Risk factors for low back pain and its relation with pain related disability and depression in a Turkish sample. *Turk Neurosurg.* 2009;19(4):327–332.
- [61] Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand.* 1983;67(6):361–370.
- [62] Aydemir O. [Validity and reliability of Turkish version of hospital anxiety and depression scale]. *Turk Psikiyatri Derg.* 1997;8(4):280–287. Turkish.
- [63] Yazici AC, Basterzi AD, Acar ST, et al. Alopesi Areata ve Aleksitimi [Alopecia areata and alexithymia]. *Turk Psikiyatri Derg.* 2006;17(2):101–106. Turkish.
- [64] Bezircioglu I, Gulseren L, Oniz A. [Depression-anxiety and disability in the premenopausal and postmenopausal period]. *Turk Psikiyatri Derg.* 2004;15(3):199–208. Turkish.
- [65] Ebrinc S, Acikel C, Basoglu C, et al. [Anxiety, depression, job satisfaction, burnout and coping with stress in nurses of a burn unit: a comparative study]. *Anadolu Psikiyatri Derg.* 2002;3(3):162–168. Turkish.
- [66] Gulseren L, Gulseren S, Hekimsoy Z, et al. [Efficacy of fluoxetine and paroxetine on depression-anxiety, quality of life, disability and metabolic control in diabetic patients with major depressive disorder: a single-blind, comparative study]. *Klin Psikofarmakol B.* 2001;11(1):1–10. Turkish.
- [67] Gulseren L, Hekimsoy Z, Gulseren S, et al. [Depression-anxiety, quality of life and disability in patients with diabetes mellitus]. *Turk Psikiyatri Derg.* 2001;12(2):89–98. Turkish.
- [68] Yazici K, Senel T, Bicer A. [Anxiety, depression and quality of life in patients with lowback pain and neck pain]. *Klinik Psikiyatri Dergisi.* 2003;6(2):95–101. Turkish.
- [69] Yazici K, Tot S, Yazici A, et al. [The association of anxiety and depression levels and sociodemographic factors in medically ill inpatients]. *New Symposium.* 2003;41(3):120–124. Turkish.
- [70] Metz CE. Basic principles of ROC analysis. *Semin Nucl Med.* 1978;8(4):283–298.
- [71] Fawcett T. An introduction to ROC analysis. *Pattern Recognit Lett.* 2006;27(8):861–874.